

1 **Revisiting the consequences of cooperative breeding**

2

3 Judith M. Burkart

4 Carel P. van Schaik

5

6 Anthropological Institute and Museum

7 University of Zurich

8

9 Corresponding author:

10 Judith Burkart

11 Anthropological Institute and Museum

12 Winterthurerstrasse 190

13 8057 Zürich

14 judith.burkart@aim.uzh.ch

15

16 Short title: Reply to Thornton & McAuliffe (2015)

1 **Abstract**

2 Thornton & McAuliffe (2015) recently questioned the hypothesis that cooperative breeding has a
3 variety of socio-cognitive consequences, invoking both logical and empirical arguments. The
4 cooperative breeding hypothesis posits that the immediate tasks associated with extensive
5 allomaternal care require motivational proximate mechanisms, such as increased social tolerance
6 or proactive prosociality which, as a side effect, also can facilitate performance in socio-
7 cognitive tasks. Eventually, over evolutionary time this constellation may also, under specific
8 conditions, facilitate increases in brain size. Thus, the CBH is not merely a modified version of
9 the social brain hypothesis, as suggested by Thornton & McAuliffe, which would posit that a
10 species has to be particularly smart in order to engage in cooperative breeding. In this reply, we
11 first clarify these conceptual issues and then systematically address their criticism of the
12 empirical evidence. We conclude that the empirical evidence for the CBH is strong in primates,
13 but that future work on lineages other than primates is required to assess its generality.

14

15

16 **Keywords**

17 Cooperative breeding hypothesis, prosociality, social tolerance, callitrichid monkeys,
18 anthropomorphism, primates, species comparisons, extensive allomaternal care

1 **Introduction**

2 In a recent issue of this journal, Thornton & McAuliffe (2015), referred to as T&M from here on,
3 questioned both the logic of, and the empirical support for, the hypothesis that cooperative
4 breeding has a variety of socio-cognitive consequences. We welcome this prominent attention to
5 the hypothesis because if it survives testing, the cooperative breeding hypothesis (CBH) will play
6 an important role in explaining why one great ape lineage became so different, and did so quite
7 rapidly, from the other great ape lineages and gave rise to our own species (Hrdy 1999).

8 According to the CBH, the proximate motivational mechanisms needed to make the
9 cooperative breeding system work smoothly under naturalistic conditions also happen to
10 facilitate cognitive *performance* (though not necessarily intrinsic *ability* per se) in a variety of
11 socio-cognitive tasks (Figure 1). This idea was inspired by the seminal work of Snowdon (2001),
12 and elaborated by Burkart & van Schaik (2010), referred to as B&vS from here on, mainly
13 focussing on primates. In particular, the CBH argues that the presence of social tolerance,
14 proactive prosociality and other motivations (see below) translates into increased performance in
15 socio-cognitive tasks of social learning, information donation and teaching, gaze understanding
16 and cooperative problem solving (Figure 1; see also Burkart 2009; Burkart & van Schaik 2010;
17 Burkart & Finkenwirth 2014). Below, we examine the evidence for these predictions in light of
18 T&M's critique.

19 Before doing so, it is important to also stress what the CBH does not claim. It does not
20 claim that a species requires particularly powerful cognitive abilities and a correspondingly big
21 brain to engage in cooperative breeding. Rather, the cognitive consequences arise because the
22 available cognitive potential is expressed differently, and in the above-mentioned contexts, more
23 efficiently (e.g. Burkart and Finkenwirth 2014, page 9). Therefore, CBH is not merely a variant

1 of the social brain hypothesis, as argued by T&M, which would predict that cooperative breeding
2 required distinct cognitive abilities. Hence, we can only agree with T&M's conclusions that
3 "*there is no compelling evidence that these processes are either unique to cooperative breeders*
4 *or particularly cognitively demanding*" (page 12) or "*that there is little evidence to suggest that*
5 *cooperative breeding entails distinct cognitive challenges*" (page 13), but want to stress that
6 these conclusions in no way contradict the CBH.

7 Once a species has been a cooperative breeder for some time, however, the greater
8 efficiency of social learning relative to independent breeders may lead to natural selection on
9 enhanced cognitive ability, and thus bigger brains. We discuss these evolutionary consequences
10 in detail below.

11

12 How to test the CBHs predictions?

13 Most of an organism's traits are affected by multiple selective forces. The CBH is
14 therefore a *ceteris paribus* hypothesis, and to test it (or indeed any other evolutionary hypothesis)
15 one cannot simply compare a random set of cooperative breeders with a random set of other
16 species. The fact that species other than cooperative breeders learn socially, for instance, is on its
17 own uninformative (see T&M, page 16). Rather, systematic comparisons that control for
18 phylogenetic relatedness and as many other factors as possible are required, such as broad
19 comparative analyses or targeted comparisons.

20 In broad comparative analyses, the aim is to identify to what extent the intensity of
21 allomaternal care¹ predicts variation in a given trait, e.g. proactive prosociality, across a large

¹ Cooperative breeding, where helpers do not breed, is toward the upper end of the allomaternal care continuum (with eusociality at its extreme, see also Isler & van Schaik 2012). Note that this continuous variable allows for powerful comparative analyses also within primates because this taxon shows considerable variation in allomaternal care.

1 number of species, by controlling for phylogenetic relatedness and other possible confounding
2 factors, such as brain size. Broad comparative analyses require large bodies of comparative data.
3 In their absence, the method of targeted comparisons can be applied (B&vS, see also SI). Here,
4 the performance of particular cooperatively breeding species is compared with that of their
5 closest, independently breeding sister taxa. Within non-human primates, this implies comparing
6 cooperatively breeding callitrichid monkeys with capuchin and squirrel monkeys. B&vS did this
7 by systematically reviewing the literature on cognitive performance in these species. Targeted
8 comparisons are less powerful statistically than broad comparisons, but when a difference is
9 found in one direction, whereas the major confounding factors would predict the opposite
10 direction, they can be highly informative. The major confounding factor when comparing
11 cognitive performance is brain size, and because capuchin and squirrel monkeys have
12 substantially larger brains than callitrichids, the targeted comparison in B&vS is biased against
13 callitrichid monkeys and thus renders this test of the CBH conservative.

14

15 **Immediate consequences: Proximate mechanisms and performance in socio-** 16 **cognitive tasks**

17 B&vS predicted that in primates, the motivational changes needed to sustain cooperative
18 breeding would include increased social tolerance, social monitoring in relaxed contexts, and
19 proactive prosociality. These proximate mechanisms, in turn, would facilitate performance in
20 social learning tasks, information donation and teaching, gaze understanding and cooperative
21 problem solving, and also pave the way for richer and more flexible vocal communication. We
22 tested these predictions with broad comparative analyses and targeted comparisons and found
23 they were met. Here we examine these predictions and T&M's criticisms of them.

1 Proximate mechanisms regulating caretaking behaviours

2

3 *Social tolerance*

4 T&M (page 15) claim that “*it is not clear whether cooperative breeders are generally*
5 *more socially tolerant than other animals, as data for systematic comparisons are lacking*”.

6 However, Burkart et al. (2014) had made such a systematic comparison and found a strong effect
7 of allomaternal care on social tolerance, as well as an effect of pair-bonding, in a set of 24 groups
8 of 15 primate species (Figure 2). Furthermore, in this data set, variation in social tolerance within
9 the callitrichid monkeys was also linked to variation in allomaternal care (Burkart 2015).
10 Importantly, the measure used by Burkart et al. (2014) is correlated with alternative measures of
11 social tolerance (Cronin *et al.* 2013). This work shows that B&vS’s prediction is clearly met in
12 primates.

13 This result may be somewhat surprising given that cooperative breeding systems are
14 often described as despotic, especially where stress-related suppression plays a role in the
15 reproductive failure of subordinates (meerkats: Young *et al.* 2006). When viewed from the
16 perspective of reproductive skew, they are indeed despotic, but the absence of reproduction by
17 many of the group’s adults need not be brought about by permanent social stress. In fact,
18 subordinates in cooperative breeders typically show lower, rather than higher stress levels
19 compared to dominants (Creel 2001), and breeders and helpers in callitrichid monkeys show
20 highly tolerant social relationships (Aureli & Schaffner 2006).

21

22

23

1 *Proactive prosociality*

2 Among primates in nature, cooperatively breeding callitrichid monkeys share
3 significantly more food than independently breeding species, and do so in qualitatively different
4 ways: not only do they allow immatures to take their food, they also offer it in a proactive (i.e.
5 spontaneous, unsolicited) way and call them in when they have found a food item (Brown,
6 Almond & van Bergen 2004; Jaeggi, Burkart & van Schaik 2010). For instance, an analysis of
7 over 7000 food sharing events in captive marmosets revealed that during peak provisioning
8 periods, adults share more than half of all pieces of food they possess, and more than one third of
9 these transfers occur proactively; roughly 10% of all shared food items are shared with other
10 adults (Martins & Burkart 2013).

11 To identify the motivational processes underlying naturalistic prosocial behaviours such
12 as food sharing, experimental studies have been conducted. As also bemoaned by T&M,
13 however, most studies are not comparable (Silk & House 2011). In fact, methodological
14 differences demonstrably affect the results of prosociality studies (Burkart & Rueth 2013). We
15 therefore decided to use the same, cognitively undemanding paradigm and identical procedures
16 to test one of the major predictions of the CBH in 24 groups of 15 primate species: that more
17 allomaternal care correlates with a greater tendency toward proactive prosociality. These data
18 strongly supported the CBH, and none of various alternative hypotheses (Burkart et al. 2014).
19 T&M raise several methodological issues with this study, and are worried by the variation within
20 callitrichids. A detailed response to the methodological issues can be found in the SI, which
21 shows the conclusions are robust. The variability within callitrichid monkeys had us puzzled as
22 well. However, it turns out that the same factor that explains variation in prosociality and

1 tolerance across primate species, i.e. allomaternal care, is also linked to variation *within*
2 callitrichid monkeys (Burkart 2015).

3

4 Cognitive consequences of cooperative breeding?

5 The studies reviewed above showed that, as predicted, proactive prosociality and social
6 tolerance increase alongside with increasing levels of allomaternal care in primates. But do these
7 proximate motivational mechanisms of allomaternal care also facilitate performance in a variety
8 of socio-cognitive tasks? The literature reviewed in B&vS suggested it does, but a general
9 concern of T&M was that these findings may be a spurious effect of sampling bias: because
10 cooperative breeders would be more tractable as study systems, they would generally perform
11 better on whichever tests one subjects them to. While one may question this assumption, in
12 particular in primates, it produces a simple prediction: we would expect increased performance
13 by cooperative breeders in both social and non-social cognitive tasks. This, however, was not the
14 case: increased performance by callitrichids was strikingly limited to the social domain.

15

16 *Social learning*

17 It has been recognized for a long time that social tolerance facilitates social learning (Coussi-
18 Korb & Fragaszy 1995). A straightforward prediction was therefore that species with extensive
19 allomaternal care engage more in social learning, and B&vS found support for this prediction.
20 For instance, Moscovice & Snowdon (2006) report that cotton-top tamarins showed very rapid
21 social learning without receiving food rewards. In contrast to capuchin monkeys, no evidence for
22 aggression or monopolization of the foraging task was found. Likewise, in directly comparable
23 tests, capuchin monkeys did not socially learn to avoid food made unpalatable with pepper

1 (Visalberghi & Addessi 2000), whereas callitrichids rapidly did so (Snowdon & Boe 2003). In
2 fact, callitrichids who had sampled the food would utter alarm calls at a high rate.

3 The review of social learning studies by Custance et al. (2002) shows the same pattern.
4 T&M challenge this conclusion, but their alternative measure does not alter the outcome:
5 cooperative breeders were still twice as likely to be successful social learners than expected.
6 Nonetheless, we agree that future surveys should ideally control for major differences in the
7 mechanisms of social learning.

8 T&M argue we should look at evidence for social learning from the wild. Evidence for
9 social learning in callitrichids in the wild is clearly available, and by no means restricted to
10 parent-offspring contexts (Gunhold, Whiten & Bugnyar 2014; Gunhold *et al.* 2014). However,
11 even though establishing social learning experimentally in the wild is possible (Reader & Biro
12 2010), not all species are equally amenable to field experiments, producing biases. Moreover,
13 identifying the actual mechanisms involved is notoriously difficult (Kendal et al. 2010). Thus,
14 wild and captive data complement each other: wild data inform us about the incidence and
15 function, whereas captive data remain indispensable to reveal the mechanism.

16 T&M also argue that other species are capable of social learning, and that many
17 cooperative breeders do not imitate. Again, we never predicted that only cooperative breeders
18 would be able to learn socially, but merely that *ceteris paribus* they are more likely to do so. In
19 fact, Burkart & Finkenwirth (2014, page 8) argued that the currently available evidence
20 “suggests that the basic underlying mechanisms for social learning, including imitation, are
21 present in many species but applied in species-specific ways”. In primates, we find a bias
22 towards cooperative breeders, with their higher social tolerance and reduced gaze aversion (De
23 Wandelaer & Burkart 2008; Burkart 2009).

1 *Teaching*

2 Another prediction by B&vS was that the readiness to share food may extend to a
3 tendency to share information, and thus to teach. T&M incorrectly assume that we implied the
4 prediction to concern intentional teaching, which we did not. They also conclude that teaching is
5 not found predominantly among cooperative breeders. This is surprising in light of Thornton &
6 Raihani's (2010, page 299) conclusion that "...to date, the strongest evidence for teaching has
7 been found in cooperatively breeding species". T&M now suggest this pattern "*may well reflect*
8 *their tractability as study systems*". Although this may hold for some lineages, it clearly does not
9 in nonhuman primates, where extensive attempts have been made to identify teaching, in
10 particular in our closest relatives, the independently breeding chimpanzees, with very limited
11 success (Moore 2013, Kline 2015).

12 Research into the taxonomic distribution of teaching, and thus resolution of this issue, is
13 hampered by debates about its operational definition. All agree, however, that teaching by adults
14 is most likely where individual learning is costly or opportunities to do so are lacking (Kline
15 2015). Teaching by felid mothers is clearly consistent with this. The same argument also predicts
16 teaching in tool-using great apes, whose immatures take years to learn how to use tools
17 (Meulman, Seed & Mann 2013). Yet, mothers do not teach, even if they could easily scaffold
18 their infants' attempts (Völter, Rossano & Call 2015). In contrast, teaching-like behaviour in
19 callitrichids is increasingly demonstrated (Dell'MourRange & Huber 2009; Rapaport 2011,
20 Chow et al. 2015, Takahashi et al. 2015), including, surprisingly, indications that they may
21 actually perceive knowledge or skill deficits in naïve individuals, implying they show intentional
22 teaching (Humble & Snowdon 2008, Moura, Nunes & Langguth 2010, Martins & Burkart 2013).

23

1 *Gaze following, cooperative problem-solving, and communication*

2 The CBH was in part inspired by the finding that common marmosets would readily
3 solve perspective-taking tasks (Burkart & Heschl 2007), which had been used as evidence for
4 simple Theory of Mind understanding, and which chimpanzees passed (Hare *et al.* 2000),
5 whereas capuchin monkeys did not (Hare *et al.* 2003). Likewise, in directly comparable
6 cooperative problem solving tasks, callitrichids cooperate at much higher rates (i.e. 97% when
7 both were rewarded, and 76% when only one cooperation partner was) than independently
8 breeding capuchin monkeys (i.e. 76% and 39%, Snowdon & Cronin 2007).

9 Especially sophisticated among callitrichid monkeys is their vocal communication
10 (Zuberbühler 2011; Snowdon 2001; Snowdon 2013; Borjon & Ghazanfar 2014, Chow *et al.*
11 2015; Takahashi *et al.* 2015), and in fact we predict that the most pervasive differences between
12 cooperative and independent breeders may well be found in communicative contexts (see also
13 Manser *et al.* 2014). Our literature review (B&vS) suggested that these are systematic patterns,
14 even though the scarcity of sufficiently comparable data from different species made this
15 conclusion preliminary.

16

17 What about other lineages?

18 The CBH, first introduced by Hrdy in 1999, argued that an ape with the life history attributes of
19 *Homo sapiens* (in particular the co-occurrence of prolonged periods of dependency combined
20 with short birth intervals) could not have evolved unless mothers received allomaternal
21 assistance. Subsequent efforts were aimed at identifying whether there was also a cognitive
22 dimension to the CBH, from the perspective of both immatures (Hrdy 2009, 2013) and helpers
23 (Burkart & van Schaik 2010, Burkart *et al.* 2009). It was thus specifically developed and

1 elaborated for the primate lineage (see also SI), and accordingly tests have so far focused on this
2 lineage. Nonetheless, whether the CBH also applies to other lineages is a valid question.

3 Phylogeny matters in two different ways. First, the predictions of the immediate
4 motivational and cognitive consequences of cooperative breeding are directly derived from the
5 specific helping behaviours performed by helpers (see Figure 1). For instance, fish helpers may
6 fan eggs or defend the territory (Taborsky & Limberger 1981), bird helpers may provision
7 offspring or mob predators (Koenig & Dickinson 2004), and primate helpers may carry the
8 young and do some provisioning (Goldizen 1987). Thus, egg fanning is unlikely to require the
9 same proximate motivational mechanisms as food sharing.

10 Second, phylogeny may affect the consequences for other aspects of behaviour. Thus, a
11 motivational change towards more prosocial attitudes to fellow group members, which is likely a
12 common trait associated with cooperative breeding across lineages, may well produce different
13 effects in a small mammal than in a chimpanzee-like, large-brained animal (Burkart et al. 2009).
14 For instance, if a great ape capable of simple Theory of Mind understanding starts to engage in
15 teaching, it is more likely to also engage in intentional teaching than a species that teaches but
16 has no Theory of Mind abilities (such as ants). This may have been a critical interaction effect in
17 human evolution (Burkart et al. 2009).

18 The extent to which the predictions of the CBH developed for primates also apply to
19 other lineages may thus critically depend on lineage-specific idiosyncrasies. In B&vS we
20 addressed this issue and showed that elephants and canids show a similar pattern: they perform
21 surprisingly well in socio-cognitive tasks but not in tasks from the non-social, physical domain.
22 But obviously more systematic comparisons are needed, along with detailed theoretical analyses
23 focusing on the interplay between new traits related to cooperative breeding (e.g. proactive

1 prosociality) and the phylogenetic substrate upon which they were grafted. Nevertheless, some of
2 the patterns identified in primates seem to even hold in phylogenetically remote lineages such as
3 eusocial insects, which offer spectacular cases of prosocial, declarative communication (bee
4 dance: Couvillon 2012), teaching (tandem running ants: Franks & Richerson 2006), social
5 learning (Leadbeater & Chittka 2007), or cognitive performance suggestive of metacognitive
6 abilities (Perry & Barron 2013) . Eusociality, represents an extreme form of cooperative
7 breeding.

8 The predictions of the CBH regarding delayed evolutionary consequences will arguably
9 be more stable across lineages because they are only indirectly linked to the idiosyncrasies of the
10 cooperative breeding systems and mainly derived from energetic considerations, as detailed in
11 the following section.

12

13 **Evolutionary consequences**

14 The evolutionary version of the CBH explores whether allomaternal care may also play a
15 role in the evolution of larger brains. The social brain hypothesis (Dunbar & Shultz 2007) is well
16 supported for birds and mammals, but it leaves much of the variation in brain size unexplained,
17 for instance in the form of so called grade-shifts, i.e. where different lineages differ strongly in
18 brain size unrelated to differences in social demands (Holekamp et al. 2015). To understand this
19 variation, it is crucial not only to focus on the benefits of large brains, but also on their costs (van
20 Schaik et al. 2012). B&vS suggested cooperative breeding played a role in brain size evolution
21 because it influences the cost-benefit balance, not because it “*entails distinct socio-cognitive*
22 *challenges*” (T&M, p. 13).

1 Big brains consume a large proportion of the energy available to an organism, in
2 particular in immatures. Moreover, this energy needs to be constantly available, because
3 developing brains cannot be starved temporarily without permanent damage. In cooperatively
4 breeding species, this constraint on brain size can be relieved by the energy subsidies to
5 immatures by alloparents, which therefore, all other things being equal, permits the evolution of
6 bigger brains in species with allomaternal care. However, selection may also favour investing the
7 load reduction of mothers receiving allomaternal care into the production of a higher number of
8 offspring, rather than in brainier offspring. The conditions under which one pathway is more
9 likely than the other are summarized under the concepts of ‘life history filter’ and ‘energy filter’
10 (van Schaik, Isler & Burkart 2012; Isler & van Schaik 2014). Empirical support from broad
11 phylogenetic comparisons is available for both pathways: in birds and some mammals,
12 allomaternal care is linked with higher reproductive rates, in particular so in nonhuman primates,
13 and with bigger brains in mammals in general (particularly so in carnivores, but not in nonhuman
14 primates; Isler & van Schaik 2012).

15 The balance between costs and benefits of larger brains is further modulated by processes
16 related to cultural intelligence. Brains per se do not provide a fitness advantage to individuals,
17 but rather the skills that are learned ontogenetically do. Since social learning is per definition
18 more efficient than individual learning, a brain of the same size will acquire more fitness-
19 relevant skills if it does so via social rather than asocial, individual learning. The cultural
20 intelligence approach thus predicts that species able to rely more systematically on social
21 learning (for whatever reason, cooperative breeding being one among others), are more likely to
22 be able to evolve a bigger brain (van Schaik & Burkart 2011; van Schaik, Isler & Burkart 2012;

1 see also Burkart in press, for how cooperative breeding, cultural intelligence and life history
2 interact in enabling the evolution of big brains).

3

4 **Conclusions**

5 In contrast to T&M, we would argue on the basis of current evidence that the support for the
6 cooperative breeding hypothesis is actually surprisingly strong, at least for primates, the lineage
7 in which systematic comparative work has been done. Of course some predictions remain
8 untested. Such tests should ideally be along the lines of Burkart et al. (2014), thus involving, as
9 called for by T&M, “*phylogenetically controlled analyses, accounting for ecological variables*
10 *such as diet and using cognitive data obtained using comparable methods*”. In particular, we
11 hope that T&M’s critique will stimulate similar comparative studies in the taxa on which T&M
12 are experts, so we can test the generality of the effects of cooperative breeding on the
13 motivational and cognitive aspects of behaviour. In biology, phylogeny matters, as do all manner
14 of confounding variables, and it is conceivable that effects are to some extent taxon-specific.
15 Mapping these effects in some detail is laborious but unavoidable.

16

17 **Acknowledgements**

18 We thank Sarah Hrdy, Karin Isler, Michèle Schubiger, and Charles Snowdon for useful
19 comments on a previous draft of this manuscript. This work was supported by the Swiss National
20 Science Foundation (SNF grants 105312-114107 and 310030-13083 to JB).

21

1 **References**

- 2 Aureli, F. & C. Schaffner, (2006) Causes, consequences and mechanisms of reconciliation: the
3 role of cooperation. In: *Cooperation in primates and humans*: 121. P. M. Kappeler & C. P.
4 van Schaik (Eds.). Springer, Heidelberg.
- 5 Borjon, J.I. & A.A. Ghazanfar (2014) Convergent evolution of vocal cooperation without
6 convergent evolution of brain size. *Brain Behav. Evol.*, **84**, 93-102.
- 7 Brown, G.R., R.E.A. Almond & Y. van Bergen (2004) Begging, stealing and offering: food
8 transfer in non-human primates. *Adv. Study Behav.*, **34**, 265-295.
- 9 Burkart, J.M., (2009) Socio-cognitive abilities and cooperative breeding. In: *Learning from*
10 *Animals? Examining the Nature of Human Uniqueness*: 123-141. L. S. Röska-Hardy & E.
11 M. Neumann-Held (Eds.). Psychology Press.
- 12 Burkart, J.M. (2015) Opposite effects of male and female helpers on social tolerance and
13 proactive prosociality in callitrichid family groups. *Sci. Rep.*, **5**, 9622.
- 14 Burkart, J.M. (in press) The evolution and consequences of sociality. In: *APA Handbook of*
15 *Comparative Psychology* (eds: Call, J. et al.)
- 16 Burkart, J.M., O. Allon, F. Amici, C. Fichtel, C. Finkenwirth, A. Heschl, J. Huber, K. Isler, Z.K.
17 Kosonen, E. Martins, E.J.M. Meulman, R. Richiger, K. Rueth, B. Spillmann, S.
18 Wiesendanger & C. P. van Schaik (2014) The evolutionary origin of human hyper-
19 cooperation. *Nature Comm.*, **5**, 4747.
- 20 Burkart, J.M. & C. Finkenwirth (2014) Marmosets as model species in neuroscience and
21 evolutionary anthropology. *Neurosci. Res.*, **93**, 8-19.
- 22 Burkart, J.M. & A. Heschl (2007) Understanding visual access in common marmosets, *Callithrix*
23 *jacchus*: perspective taking or behaviour reading? *Anim. Behav.*, **73**, 457-469.
- 24 Burkart, J.M., Hrdy, S.B., & van Schaik, C.P. (2009). Cooperative breeding and human cognitive
25 evolution. *Evol. Anthropol.*, *18*(5), 175-186.
- 26 Burkart, J.M. & K. Rueth (2013) Preschool children fail primate prosocial game because of
27 attentional task demands. *PLoS ONE*, **8**, e68440.
- 28 Burkart, J.M. & C.P. van Schaik (2010) Cognitive consequences of cooperative breeding in
29 primates. *Anim. Cogn.*, **13**, 1-19.
- 30 Chow, C.P., Mitchell, J.F., & Miller, C.T. (2015). Vocal turn-taking in a non-human primate is
31 learned during ontogeny. *Proc. Roy. Soc. B*, *282*(1807), 20150069.
- 32 Coussi-Korbel, S. & D.M. Fragaszy (1995) On the relation between social dynamics and social
33 learning. *Anim. Behav.*, **50**, 1441-1453.
- 34 Couvillon M (2012) The dance legacy of Karl von Frisch. *Insects Soc.* *59*:297–306
- 35 Creel, S. (2001) Social dominance and stress hormones. *Trends Ecol. Evol.*, **16**, 491-497.
- 36 Cronin, K.A., E.J. van Leeuwen & D. Haun, (2013) The many metrics of social tolerance. *Folia*
37 *Primatol.*, **84**, 261-262.
- 38 Custance, D. M., A. Whiten & T. Fredman (2002) Social learning and primate reintroduction.
39 *Int. J. Primatol.*, **23**, 479-499.
- 40 De Wandelaer, T. & J.M. Burkart (2008) Is there a relationship between gaze aversion and social
41 structure? *Folia Primatol.*, **79**, 396-397.
- 42 Dell'Mour, V., F. Range & L. Huber (2009) Social learning and mother's behavior in
43 manipulative tasks in infant marmosets. *Am. J. Primatol.*, **71**, 503-509.
- 44 Franks, N. R., & Richardson, T. (2006) Teaching in tandem-running ants. *Nature*, *439*(7073),
45 153-153.

- 1 Goldizen, A.W., (1987) Tamarins and marmosets: Communal care of offspring. In: *Primate*
 2 *Societies*: 34-43. Smuts. B. et al. (Ed.). The University of Chicago Press, Chicago.
- 3 Gunhold, T., J.M. Massen, N. Schiel, A. Souto & T. Bugnyar (2014) Memory, transmission and
 4 persistence of alternative foraging techniques in wild common marmosets. *Anim. Behav.*,
 5 **91**, 79-91.
- 6 Gunhold, T., A. Whiten & T. Bugnyar (2014) Video demonstrations seed alternative problem-
 7 solving techniques in wild common marmosets. *Biol. Lett.*, **10**, 20140439.
- 8 Hare, B., E. Addessi, J. Call, M. Tomasello & E. Visalberghi (2003) Do capuchin monkeys know
 9 what conspecifics do and do not see *Anim. Behav.*, **65**, 131-142.
- 10 Hare, B., J. Call, B. Agnetta & M. Tomasello (2000) Chimpanzees know what conspecifics do
 11 and do not see. *Anim. Behav.*, **59**, 771-785.
- 12 Holekamp, K.E., Dantzer, B., Stricker, G., Yoshida, K.C.S., & Benson-Amram, S. (2015).
 13 Brains, brawn and sociality: a hyaena's tale. *Animal Behaviour*, **103**, 237-248.
- 14 Humle, T. & C.T. Snowdon (2008) Socially biased learning in the acquisition of a complex
 15 foraging task in juvenile cottontop tamarins. *Anim. Behav.*, **27**, 267-277.
- 16 Hrdy, S. (1999) *Mother Nature*. New York: Pantheon Books.
- 17 Hrdy, S. (2009) *Mothers & Others*. Cambridge: Harvard University Press.
- 18 Hrdy, S. (2013) Development + social selection in the emergence of “emotionally modern”
 19 humans. *New Frontiers in Social Neuroscience*. Springer International Publishing, 2014.
 20 S. 57-91.
- 21 Isler, K. & C.P. van Schaik (2012) Allomaternal care, life history and brain size evolution in
 22 mammals. *J. Hum. Evol.*, **63**, 52-63.
- 23 Isler, K. & C.P. van Schaik (2014) How humans evolved large brains: Comparative evidence.
 24 *Evol. Anthropol.*, **23**, 65-75.
- 25 Jaeggi, A., J.M. Burkart & C.P. van Schaik (2010) On the psychology of cooperation in humans
 26 and other primates: The natural history of food sharing and experimental evidence of
 27 prosociality. *Phil. T. Roy. Soc. B.*, **12**, 2723-2735.
- 28 Kendal, R.L., Galef, B.G., & Van Schaik, C.P. (2010) Social learning research outside the
 29 laboratory: How and why?. *Learning & Behavior*, **38**, 187-194.
- 30 Kline, M.A. (2015). How to learn about teaching. *Behav. Brain Sci.*, **38**, 1-70.
- 31 Koenig, A. & J. Dickinson (2004) *Ecology and Evolution of Cooperative Breeding in Birds*.
 32 Cambridge: Cambridge University Press.
- 33 Leadbeater, E., & Chittka, L. (2007) Social learning in insects—from miniature brains to
 34 consensus building. *Current Biology*, 17(16), R703-R713.
- 35 Manser, M. B., Jansen, D, Graw, B., Hollén, L. I., Bousquet, C. A., Furrer, R. D., & le Roux, A.
 36 (2014). Vocal complexity in meerkats and other mongoose species. *Adv. Stud. Behav.*,
 37 **46**, 281-310.
- 38 Martins, E.M.G. & J.M. Burkart (2013) Common marmosets preferentially share difficult to
 39 obtain food items. *Folia Primatol.*, **84**, 281-282.
- 40 Meulman, E J.M., Seed, A.M., & Mann, J. (2013). If at first you don't succeed... Studies of
 41 ontogeny shed light on the cognitive demands of habitual tool use. *Phil Trans B*, **368**,
 42 20130050.
- 43 Moore, R. (2013) Social learning and teaching in chimpanzees. *Biol. Phil.*, **28**, 879-901.
- 44 Moscovice L.R., & Snowdon C.T. (2006) The role of social context and individual experience in
 45 novel task acquisition in cottontop tamarins.. *Anim. Behav.* 71:933-943.


- 1 Moura, A.C., H.G. Nunes & A. Langguth (2010) Food sharing in lion tamarins: Does foraging
2 difficulty affect investment in young by breeders and helpers? *Int. J. Primatol.*, **31**, 848-
3 862.
- 4 Perry, C. J., & Barron, A. B. (2013) Honey bees selectively avoid difficult choices. *PNAS*
5 110(47), 19155-19159.
- 6 Rapaport, L.G. (2011) Progressive parenting behavior in wild golden lion tamarins. *Behav. Ecol.*,
7 **22**, 745-754.
- 8 Reader, S.M., & Biro, D. (2010). Experimental identification of social learning in wild animals.
9 *Learn. Behav.*, **38**(3), 265-283.
- 10 Silk, J. B. & B. R. House (2011) Evolutionary foundations of human prosocial sentiments.
11 *PNAS*, **108**, 10910-10917.
- 12 Snowdon, C.T., (2013) Language parallels in New World primates. In: *Animal Models of Speech*
13 *Lang. Disord.*: 241-261. Springer, New York.
- 14 Snowdon, C.T. (2001) Social processes in communication and cognition in callitrichid monkeys:
15 a review. *Anim. Cogn.*, **4**, 247-257.
- 16 Snowdon C.T., and Boe C.Y. (2003). Social communication about unpalatable foods in tamarins.
17 *J. Comp. Psychol.* 117(2):142-148.
- 18 Snowdon, C.T., & Cronin, K.A (2007) Cooperative breeders do cooperate. *Behav. Proc.* **76**,138-
19 141.
- 20 Taborsky, M. & D. Limberger (1981) Helpers in fish. *Behav. Ecol. Sociobiol.*, **8**, 143-145.
- 21 Takahashi, D.Y., Fenley, A.R., Teramoto, Y., Narayanan, D.Z., Borjon, J. I., Holmes, P., &
22 Ghazanfar, A.A. (2015). The developmental dynamics of marmoset monkey vocal
23 production. *Science*, 349(6249), 734-738.
- 24 Thornton, A., & Raihani, N.J. (2010). Identifying teaching in wild animals. *Learning &*
25 *Behavior*, **38**(3), 297-309.
- 26 Thornton, A. & K. McAuliffe (2015) Cognitive consequences of cooperative breeding? A critical
27 appraisal. *J. Zool.*, **295**, 12-22.
- 28 van Schaik, C.P. & J.M. Burkart (2011) Social learning and evolution: the cultural intelligence
29 hypothesis. *Phil. T. Roy. Soc. B.*, **366**, 1008-1016.
- 30 van Schaik, C.P., K. Isler & J.M. Burkart (2012) Explaining brain size variation: from social to
31 cultural brain. *Trends Cogn. Sci.* **16**, 277-284
- 32 Visalberghi E., and Addessi E. (2000). Response to changes in food palatability in tufted
33 capuchin monkeys. *Anim. Behav.* 59:231-238.
- 34 Völter, C., F. Rossano & J. Call (2015) From exploitation to cooperation: social tool use in
35 orang-utan mother-offspring dyads. *Anim. Behav.*, **100**, 126-134.
- 36 Young, A.J., A.A. Carlson, S.L. Monfort, A.F. Russell, N.C. Bennett & T. Clutton-Brock (2006)
37 Stress and the suppression of subordinate reproduction in cooperatively breeding
38 meerkats. *PNAS*, **103**, 12005-12010.
- 39 Zuberbühler, K., (2011) Cooperative breeding and the evolution of vocal flexibility. In: *The*
40 *Oxford Handbook of Language Evolution*. M. Tallerman & K. Gibson (Eds.).

41
42

1 **Figure Legends**

2

3



4

5 **Figure 1**

6 The hypothesized pathway for a link between cooperative breeding and socio-cognitive

7 performance. The CBH predicts that cooperative breeding – or extensive allomaternal care – is

8 associated with a set of proximate mechanisms necessary to support and enable allomaternal


9 behaviors. At the same time, these proximate mechanisms facilitate performance (but not

10 necessarily ability) in a variety of socio-cognitive tasks. For instance, cooperatively breeding

11 primates show increased levels of social tolerance, which is necessary to ensure smooth infant

12 transfers in the canopy At the same time, however, social tolerance also facilitates performance

13 in social learning tasks (see text).



1

2 **Figure 2**

3 Relationship between allomaternal care and social tolerance in primates. Light green dots: pair

4 bonded species; blue circles: species qualifying as cooperative breeders.

5


1 Supplementary Information

2

3 1) Targeted comparisons

4 When broad comparative analyses are not possible to test evolutionary hypotheses, targeted
 5 comparisons can be performed, i.e. comparisons between closely related sister taxa that differ
 6 with regard to the relevant factor but are as similar as possible with regard to potential
 7 confounding factors. The necessity to focus on targeted comparisons in closely related species,
 8 rather than comparing any species that differ with regard to the relevant factor is depicted in
 9 Figure 2. This figure exemplifies how the comparison of randomly chosen species, one an
 10 independent breeder and one a cooperative breeder, may result in the independent breeder
 11 outperforming the cooperative breeder even if in reality, the trait in question is indeed positively
 12 linked with cooperative breeding. It is thus important to keep in mind that the comparison of any
 13 two randomly chosen species, one breeding cooperatively but the other one breeding
 14 independently, has no explanatory power for the CBH if we cannot exclude other factors that
 15 may also affect the trait of interest.

16



17

18 Figure 2

19 Schematic depiction of a situation where cooperative breeding has a positive influence on cognitive performance
 20 (green dotted regression lines: cooperative breeders; red dashed line: independent breeders; c_1 = cooperatively
 21 breeding species 1, i_1 = independently breeding sister taxon of c_1 ; etc.). Targeted comparisons show that in each
 22 species pair, the cooperatively breeding species outperforms its sister taxon. However, if additional factors also play
 23 a role (factor x, e.g. brain size), comparing non-sister taxa that vary widely with regard to this factor is
 24 uninformative or may even wrongly lead to the opposite conclusion (because i_2 outperforms c_1 by distance d).

25

1 **2) Methodological issues of the group service study (Burkart et al. 2014)**

2 In the group service paradigm, which had been designed to provide a comparable
 3 measure of proactive prosociality across species, one animal can pull food toward the group but
 4 cannot get any of it for itself. The apparatus is very simple and prior to the test, the animals have
 5 to demonstrate they understand it; during the test, several control conditions are included to
 6 make sure they continue to do so. The paradigm had been piloted with three species and found to
 7 be robust (Burkart & van Schaik 2013). The subsequent comparative study, involving 24 groups
 8 of 15 different primate species, found strong support for the CBH, and none for a range of other
 9 hypotheses. Thornton & McAuliffe (2015) raise three issues with the group service, (i) that it is
 10 limited to primates only, (ii) that there is substantial variation within callitrichids, and (iii)
 11 methodological issues. While (ii) has been addressed empirically (Burkart et al. 2015) as well as
 12 in the main text, (i) and (iii) are discussed below.

13

14 *Re (i): Main focus on primates and anthropocentrism*

15 T&M criticize our approach as anthropocentric. In one sense, we accept this criticism.
 16 The aim of evolutionary anthropology is to try and understand why humans became so different
 17 from the other great apes. We hypothesise that extensive allomaternal care was partly, perhaps
 18 even largely responsible for this. In other words, we assume that a non-unique process,
 19 convergently found in other lineages, has had predictable effects in our own lineage. Our main
 20 comparative effort focused on primates for this reason as well. In fact, the CBH developed by
 21 Hrdy in 1999 was that an ape with the life history attributes of *Homo sapiens* (in particular the
 22 co-occurrence of prolonged periods of dependency combined with short birth intervals) could not
 23 have evolved unless mothers had allomaternal assistance. Subsequent efforts were aimed at
 24 identifying whether there was also a cognitive dimension to the CBH, both from the perspective
 25 of helpers (Burkart & van Schaik 2010, Burkart et al. 2009) and of immatures (Hrdy 2009,
 26 2013). Neither version requires actors involved to be particularly brainy to start with.

27 The charge of anthropocentrism is not accurate when read in the sense that we project
 28 human abilities onto the animals or are only interested in those features that have direct human
 29 equivalents. We apply the comparative method and aim to understand the evolution of our own
 30 species as one would try to understand the evolution of any other species using this approach.
 31 Moreover, primates are not a bad lineage in which to examine the consequences of allomaternal

1 care because they show quite pronounced variation in this feature, in fact much more so than any
2 other mammal lineage (see Figure 1 in Isler & van Schaik 2012).

3
4

5 *Re (iii): Methodological issues*

6 T&M are concerned that some results were artificially produced by order effects because
7 the experiment was run in phases. In phase III of the experiment, we trained all individuals to
8 understand the apparatus, i.e. they had to be able to pull the handle with one hand, and
9 simultaneously take a piece of food with the second hand, in a position as remote as possible so
10 that they would still be able to just reach the food when completely extending their arms. They
11 thus learned that the board would pull back automatically if it was released, and had to pass
12 predefined criteria to proceed to the next phase. In phase IV of the experiment, in which we
13 assessed prosociality, we ran 5 test sessions and 5 control sessions on alternating days, each
14 consisting of a total of 70 trials. In test sessions, the board was baited with a piece of food in a
15 position which made it impossible for an individual to pull the handle and retrieve the food by
16 itself at the same time. Food could only be made available to the group if one individual would
17 pull the board and another would retrieve the food. In control sessions, the board was not baited
18 but the empty food bowl was audibly manipulated, to control for the possibility that individuals
19 would pull for reasons other than providing food. Only individuals who pulled more in test vs.
20 control sessions were included in the analyses. Furthermore, T&M wrongly criticized that we
21 hadn't used difference scores as dependent variables, but in fact we had done so (Burkart et al.
22 2014, Supplementary Information, page 22), but this did not affect the conclusion. Finally, in
23 both test and control sessions, so called motivation trials were interspersed, identical to phase III,
24 where food could be made available individually. In the motivation trials, food items were
25 virtually always taken, which clearly demonstrates that over test session, the subjects had not
26 forgotten how the apparatus worked.

27 For those species that had shown consistent provisioning over all test sessions in phase
28 IV, we had added an additional control condition in phase V to exclude further alternatives for
29 high levels of provisioning. Phase V was identical to phase IV, but access to the food in the more
30 distant position was always blocked (but still available in motivation trials). T&M argued that
31 the animals pulled less in phase V, not because they understood that they could no longer
32 provide food, but because of a general decline in pulling over the experiment. If this argument

1 were correct, the conclusion that increased allomaternal care elicits increased proactive
2 prosociality would no longer be compelling.

3 There are at least two lines of evidence against the interpretation suggested by T&M.
4 First, supplementary Figure 3 in Burkart et al. (2014) clearly shows that species with high levels
5 of allomaternal care do not reduce their pulling activity over the time of the experiment. Over the
6 test sessions in phase IV, we found three clusters. The first contained groups with provisioning
7 rates that were low throughout, the second contained groups with decreasing rates, and the third
8 groups with stable high or increasing rates of provisioning. Tellingly, species from the first
9 cluster had the lowest level of allomaternal care, the second cluster slightly higher, and the third
10 cluster had the highest levels of allomaternal care. It was this third group of species that had been
11 subjected to the final phase V, to provide additional evidence that they indeed understood the
12 consequences of their pulling (which is obviously not possible for species who do not pull for
13 others in the first place), and they passed this phase. Second, for two callitrichid groups we re-
14 ran phase IV tests after the end of the experiment, with the result that they immediately resumed
15 pulling for their group members at similar rates as before (Burkart et al. 2014, Supplementary
16 Information, page 15). In the meantime, additional callitrichid groups have been tested again
17 with phase IV trials for various reasons, and they also immediately resumed to pull for each
18 other (unpublished data). In fact, we now use the group service apparatus as an environmental
19 enrichment device for the callitrichid groups at our primate station, and it is possible to provide
20 entire meals via these devices. The published finding is therefore robust.

21 Lest this conclusion be misinterpreted, we stress that prosocial behaviour may be
22 heterogeneous in terms of its motivating factors (Jaeggi, Burkart & van Schaik 2010). Hence,
23 these results do not mean, as implied by T&M, that species without extensive allomaternal care
24 do not show prosocial behaviour. They do, but it is more commonly solicited by the proximity
25 and behaviour of a special friend or relative. Moreover, we do not exclude other sources of
26 proactive prosociality, e.g. between particular highly bonded dyads, in other species, but so far
27 there is no clear evidence that that occurs. Again, the hypothesis is a *ceteris paribus* one, like
28 virtually all ideas in biology.

29

30 **References**

- 1 Burkart, J.M., O. Allon, F. Amici, C. Fichtel, C. Finkenwirth, A. Heschl, J. Huber, K. Isler, Z.K. Kosonen, E.
2 Martins, E.J.M. Meulman, R. Richiger, K. Rueth, B. Spillmann, S. Wiesendanger & C. P. van Schaik
3 (2014) The evolutionary origin of human hyper-cooperation. *Nature Comm.*, **5**, 4747.
4 Burkart, J.M., Hrdy, S.B., & van Schaik, C.P. (2009). Cooperative breeding and human cognitive evolution. *Evol.*
5 *Anthropol.*, *18*(5), 175-186.
6 Burkart, J.M. & C.P. van Schaik (2010) Cognitive consequences of cooperative breeding in primates. *Anim. Cogn.*,
7 **13**, 1-19.
8 Burkart, J.M. & C.P. van Schaik (2013) Group service in macaques, capuchins and marmosets: A comparative
9 approach to identifying proactive prosocial motivations. *J. Comp. Psychol.*, **127**, 212-226.
10 Hrdy, S. (1999) *Mother Nature: A History of Mothers, Infants, and Natural Selection*. New York: Pantheon Books.
11 Hrdy, S. (2009) *Mothers & Others: The Evolutionary Origins of Mutual Understanding*. Cambridge: Harvard
12 University Press.
13 Hrdy, S. (2013) Development + social selection in the emergence of “emotionally modern” humans. In: *New*
14 *Frontiers in Social Neuroscience*. Springer International Publishing, 2014. S. 57-91.
15 Isler, K. & C.P. van Schaik (2012) Allomaternal care, life history and brain size evolution in mammals. *J. Hum.*
16 *Evol.*, **63**, 52-63.
17 Jaeggi, A., J.M. Burkart & C.P. van Schaik (2010) On the psychology of cooperation in humans and other primates:
18 The natural history of food sharing and experimental evidence of prosociality. *Phil. T. Roy. Soc. B.*, **12**,
19 2723-2735.
20 Thornton, A. & K. McAuliffe (2015) Cognitive consequences of cooperative breeding? A critical appraisal. *J. Zool.*,
21 **295**, 12-22.
22
23